
SHARE WORKING PAPER SERIES

Precautionary Behaviours of Individuals with Multimorbidity during the COVID-19 Pandemic

Alice Delerue Matos, Andreia F. Paiva, Cláudia Cunha, Gina Voss

Working Paper Series 69-2021

DOI: [10.17617/2.3332949](https://doi.org/10.17617/2.3332949)

SHARE-ERIC | Amalienstr. 33 | 80799 Munich | Germany | share-eric.eu



mea | MAX PLANCK INSTITUTE FOR
SOCIAL LAW AND SOCIAL POLICY
Munich Center for the Economics of Aging



This project has received funding from the European Union under grant agreements VS/2019/0332, VS/2020/0313 and the European Union's Horizon 2020 research and innovation programme under grant agreements No 870628, No 101015924.



SPONSORED BY THE
Federal Ministry
of Education
and Research

Supported by the



About the SHARE Working Paper Series

The series is designed to provide a timely discussion of results based on SHARE data within the SHARE family, i.e., members of the SHARE Country Teams, Area Coordination Teams and other SHARE bodies. The papers are not peer reviewed; the authors are solely responsible for the scientific content and the graphical layout of their submissions. The respective Country Team Leaders and Area Coordinators are encouraged to look over the submissions by their team members.

The publisher (SHARE ERIC) checks working papers in this series for formal issues such as proper acknowledgements to the funders of SHARE. The publisher takes no responsibility for the scientific content of the paper.

Acknowledgements

This publication is based on preliminary SHARE wave 8 release 0 data (Börsch-Supan 2020). Therefore, the analyses, conclusions and results are preliminary. Please see Scherpenzeel et al. (2020) for methodological details. In addition, this paper uses data from SHARE Wave 7 (DOI: 10.6103/SHARE.w7.710), see Börsch-Supan et al. (2013) for methodological details. The SHARE data collection has been funded by the European Commission, DG RTD through FP5 (QLK6-CT-2001-00360), FP6 (SHARE-I3: RII-CT-2006-062193, COMPARE: CIT5-CT-2005-028857, SHARELIFE: CIT4-CT-2006-028812), FP7 (SHARE-PREP: GA N°211909, SHARE-LEAP: GA N°227822, SHARE M4: GA N°261982, DASISH: GA N°283646) and Horizon 2020 (SHARE-DEV3: GA N°676536, SHARE-COHESION: GA N°870628, SERISS: GA N°654221, SSHOC: GA N°823782) and by DG Employment, Social Affairs & Inclusion through VS 2015/0195, VS 2016/0135, VS 2018/0285, VS 2019/0332, and VS 2020/0313. Additional funding from the German Ministry of Education and Research, the Max Planck Society for the Advancement of Science, the U.S. National Institute on Aging (U01_AG09740-13S2, P01_AG005842, P01_AG08291, P30_AG12815, R21_AG025169, Y1-AG-4553-01, IAG_BSR06-11, OGHA_04-064, HHSN271201300071C, RAG052527A) and from various national funding sources is gratefully acknowledged (see www.share-project.org).

This study was financially supported by the Portuguese Foundation for Science and Technology through national funds (PIDDAC) and co-financed by the European Regional Development Fund (ERDF), through the Operational Programme PORNorte and PORLisboa, AAC No 01/SAICT /2016, Application No 022209 - DATALAB.

PRECAUTIONARY BEHAVIOURS OF INDIVIDUALS WITH MULTIMORBIDITY
DURING THE COVID-19 PANDEMIC

Alice Delerue Matos
Department of Sociology
Communication and Society Research Centre
Institute of Social Sciences
University of Minho
Portugal
adelerue@ics.uminho.pt

Andreia F. Paiva
Communication and Society Research Centre
Institute of Social Sciences
University of Minho
Portugal
andreiafonsecapaiva@gmail.com

Cláudia Cunha
Communication and Society Research Centre
Institute of Social Sciences
University of Minho
Portugal
claudiacunha@ics.uminho.pt

Gina Voss
Communication and Society Research Centre
Institute of Social Sciences
University of Minho
Portugal
ginavoss@ics.uminho.pt

Abstract: Studies show that older individuals with multimorbidity are more susceptible to develop a more severe case of COVID-19 when infected by the virus. These individuals are more likely to be admitted to Intensive Care Units and to die from COVID-19 related conditions than younger individuals or those without multimorbidity. This research aimed to assess whether there are differences in terms of precautionary behaviours between individuals aged 50+ with multimorbidity and their counterparts without multimorbidity residing in 25 European countries plus Israel. We used data from the SHARE Corona Survey on the socio-demographic and economic characteristics, multimorbidity, and precautionary behaviours of individuals. SHARE wave 8 and 7 databases were also used to fully identify individuals with multimorbidity. Our results showed that individuals with multimorbidity were more likely to exhibit precautionary behaviours than their counterparts without multimorbidity when gender, age, education, financial distress and countries were included as controls. Additionally, we found that women, more educated individuals and those experiencing more financial distress adopt more protective behaviours than their counterparts. Our results also indicate that the prevalence of precautionary behaviours is higher in Southern Europe and lower in Northern countries. To guarantee the adoption of preventive actions against COVID-19, public health messaging and actions must continue to be disseminated among middle and older aged persons with multimorbidity, and more awareness campaigns should be targeted at men and less educated individuals but also at persons experiencing less financial distress, particularly in countries where people engaged in fewer precautionary behaviours.

Keywords: COVID-19; SHARE; Precautionary Behaviours; Multimorbidity; Pandemic; Public Health

INTRODUCTION

Emerging infectious diseases that progress rapidly from a local epidemic to a pandemic are a significant threat to global public health. The most recent example, the COVID-19 pandemic, has impelled the authorities to intervene quickly and effectively to reduce the dissemination of the virus. A critical element to limit the spread of the disease is rapid and large-scale behavioural changes. And, indeed, authorities worldwide have responded with a set of guidelines aimed at changing people's behaviours. These guidelines include vigorous hand washing, physical distancing, staying at home as much as possible, maintaining respiratory hygiene, wearing a face mask, avoiding touching the face with unclean hands and disinfecting frequently touched objects and surfaces (Ejaz et al., 2020; Shankar et al., 2020). Although these guidelines are aimed at everyone, they should be especially followed by individuals who are at greater risk of suffering complications from the disease when infected by the virus.

A growing body of evidence suggests that the risk of developing severe life-threatening complications in COVID-19 pandemic is higher, namely, in individuals who are older and/or suffering from multimorbidity (Ramage-Morin & Polsky, 2020; Sanyaolu et al., 2020; Zaninotto et al., 2020), defined by Marengoni (2011) as the co-existence of two or more chronic conditions. When infected, these individuals have an increased likelihood of hospitalization, admission to intensive care units (ICU) and even death (Boddington et al., 2020; Docherty et al., 2020; Froes et al., 2020; Guan et al., 2020; Iaccarino et al., 2020; Zhang et al., 2020).

The theoretical background on behavioural research in response to pandemics is heavily focused on theories of risk perception, such as the Health Belief Model (HBM) (Strecher et al., 1997). According to the HBM, whether or not an individual will choose to act on a health-related behaviour will build on the perception of four aspects: susceptibility to a health threat; severity of said threat; likelihood of reducing the threat by engaging in health-related behaviour, and costs or barriers that can be associated with engaging in said behaviour. HBM predicts that higher perceived threat leads to a higher likelihood of engagement in health-promoting behaviours (Glanz & Bishop, 2010; Strecher et al., 1997). In a study by Jose et al. (2021) using the HBM to characterize the perceptions and behavioural change of individuals regarding COVID-19 and control measures, the authors found that individuals who practised hand washing, measures that prevent the transmission of infection, perceive that contracting COVID-19 would be more serious and more susceptible. They also found that older adults who had comorbid conditions also reported that contracting COVID-19 would be very severe.

There are other theories of risk perception, such as the Protection Motivation Theory (PMT) (Rogers & Prentice-Dunn, 1997) that can provide useful structures to understand behavioural choices during global challenges. In a review of literature, Bish & Michie (2010) found studies that point out that factors of the PMT, such as perceived severity of a threat, perceived vulnerability to said threat and response-efficacy, were linked to protective behaviours against infectious diseases.

A common underlying idea of these theories is that, in the presence of a threat, the individuals who perceive themselves to be more vulnerable to said threat, will increasingly engage in risk-prevention behaviours (Bonem et al., 2015; Wise et al., 2020). Similarly, individuals who perceive themselves as having a lower risk of developing illness are more likely to engage in unhealthy, risky behaviours (Jose et al., 2021). Understanding theories of behaviour changes is important to implement interventions that aim at promoting healthy behaviours and improving effective public health programs (Glanz & Bishop, 2010).

Therefore, health problems, including multimorbidity, may lead to a perception of increased vulnerability to the serious repercussions of the COVID-19 disease, consequently motivating greater precautionary behaviours to avoid infection (Jose et al., 2021).

In addition to the health condition, research outcomes elucidate the relationship between individual demographic and socioeconomic characteristics, on the one hand, and precautionary behaviours during pandemics, on the other hand.

Several researchers explain that women are slightly more afraid of the infection and perceive the disease as more severe, implementing more protective behavioural changes (Cvetković et al., 2020; Kim & Crimmins, 2020; Lep et al., 2020; Lüdecke & von dem Knesebeck, 2020; Zickfeld et al., 2020). Moreover, older men are less worried about the COVID-19 pandemic and were less cautious, according to a study conducted by Barber et al (2020).

In a literature review on determinants of precautionary behaviours during pandemics, Bish and Michie (2010) conclude that the relationship between age and precautionary behaviours is not completely clear, although most studies point that increasing age is associated with a greater likelihood of adopting precautionary behaviours.

The level of education and the income, which point to the social position of the individual, are some of the characteristics that are often taken into account in the analysis of the social determinants of health behaviours.

But, regarding education, the pattern of findings in pandemic studies is not straightforward. While some studies indicate that having a higher educational level is correlated with more preventive behaviours during pandemics (Lüdecke & von dem Knesebeck, 2020; Zickfeld et al., 2020), others reveal the opposite results (Tang & Wong, 2005) or no association at all (Tang & Wong, 2003).

Income is considered in a much smaller number of studies on health behaviours in the context of a pandemic. In a research conducted in the United States of America (USA), Jay et al (2020) concluded that residents of low-income neighbourhoods were less likely to stay at home, even with state orders, due to the need to work outside the home.

We aim at describing whether the adoption of precautionary behaviours differs among individuals aged 50+ with and without multimorbidity, in 25 European countries and Israel. Inspired by the risk perception theories, the underlying hypothesis is that individuals with multimorbidity behaved more cautiously than those without multimorbidity, controlling for main demographic and socio-economic characteristics of the individuals that are associated with precautionary behaviours. Since precautionary behaviours also depends on the characteristics of the context in which individuals operate, the variability of countries is also taken into account.

This study goal is to fill a gap in knowledge about precautionary behaviours during a pandemic. Studies comparing two population groups with different health risks, such as this one involving individuals with and without multimorbidity, on the health behaviours adopted in a pandemic context are scarce. Furthermore, there is a lack of studies on Europe and comparative studies between countries. This research is, as far as we know, the first European cross-national study of precautionary behaviours during the first wave of COVID-19 pandemic, involving a very significant number of countries.

DATA AND METHODS

Study design and setting

The current study uses data from the SHARE Corona Survey (release 0) mainly, but also from wave 8 (release 0) and 7 (release 7.1.0) databases, that were used to identify individuals with multimorbidity. For more methodological details, please see Börsch-Supan (2020a, 2020b), Börsch-Supan et al (2013), Bergmann et al (2019) and Scherpenzeel et al (2020). The sample was restricted to respondents aged 50+ who answered “yes” to

the routing question “Since the outbreak of Corona, have you ever left your home?” as those were the only ones who answered all the precautionary behaviours questions. The study focuses on 25 European countries plus Israel. The Netherlands was excluded due to the lack of observations in our interest variable. Hence, the sample size for this study is 41532 individuals.

Measures

Outcome variable

The precautionary behaviours considered in this research were: (1) going shopping; (2) going out for a walk; (3) meeting more than 5 people from outside the household; (4) visiting other family members since the outbreak (answers were reclassified into two groups: About the same/More often, and Not anymore/Less often). Additional precautionary behaviours were (5) wearing a face mask in public; (6) keeping distance from others in public (answers were reclassified into two groups: Always/Often, and Sometimes/Never). Finally, some more precautionary behaviours were assessed with the yes/no questions: (7) washed hands more than usual; (8) used hand sanitiser or disinfectant fluids more than usual; and (9) covered coughs and sneezes more than usual. A person was categorized as having high precautionary behaviours if she/he reported 7 or more of the above-mentioned behaviours and having low precautionary behaviours if fewer than 7 behaviours were reported. The cut point was based on the median value of the precautionary behaviours, by country.

Interest variable

Multimorbidity was defined as reporting two or more chronic conditions (Marengoni et al., 2011). This study uses the following chronic conditions: hip fracture; diabetes or high blood sugar; high blood pressure or hypertension; heart attack including myocardial infarction or coronary thrombosis or any other heart problem including congestive heart failure; chronic lung disease such as chronic bronchitis or emphysema; cancer or malignant tumour, including leukaemia or lymphoma, but excluding minor skin cancers; another illness or health condition. These conditions were reported in the SHARE Corona questionnaire, using the questions “Since we last interviewed you, were you diagnosed with a major illness or health condition?” and “Do you have any of the following illnesses or health conditions?”. Since individuals may suffer from non-recent multimorbidity, if they answered “no” to the first question above, the same questions from wave 8 or wave 7 were used to fully classify respondents into two groups: with multimorbidity and without multimorbidity.

Control variables

The control variables in the model were selected based on the literature review. Gender and age of the respondent at the time of the interview were selected as control variables. To take into account the socioeconomic position of the respondents, two indicators were used: education and financial distress. Education was measured according to the highest level of education attained using the standardized coding of the International Standard Classification of Education (ISCED-97). This variable was categorized into three groups: low education (ISCED-97 levels 0-2 corresponding to lower secondary school at the most); medium education (ISCED-97 level 3, upper secondary school) and high education (ISCED-97 levels 4-6 corresponding to post-secondary school). Since income was not available in the SHARE Wave 8 COVID-19 data and it is an indicator that can vary substantially in a short period of time, therefore discouraging the use of information reported in previous waves, we used a proxy indicator of income, available in SHARE Corona Survey: financial distress, assessed by the question “Thinking of your household's total monthly income since the outbreak of Corona, would you say that your household is able to make ends meet?”. The answers were reclassified into two groups: ‘with great difficulty’/‘with some difficulty’, and ‘fairly easily’/‘easily’.

Statistical analyses

This study was carried out in two stages. Firstly, to characterize our study population, descriptive statistics were applied using calibrated individual weights, since the SHARE survey does not have a uniform sample design. To analyse whether there are differences between the high and low precautionary behaviour groups, statistical tests for two-group comparison were performed (t-test (t) and chi-square tests (X^2)). To complement these analyses, we used Cohen’s d/Phi effect size measure to assess the magnitude of the observed effect on our sample. Confidence Intervals (CI) for these observed effect measures were also calculated.

Secondly, to examine the association between multimorbidity and precautionary behaviours, a multilevel logistic regression, with individuals as level one and countries as level two, was performed. As a first step, the null model (Model 1) was tested as a means to determine the variance of precautionary behaviours that are explained by country differences. The Intraclass Correlation Coefficient (ICC) of the null model is 13.6 per cent, higher than the recommended cut-point of 5 per cent and, for this reason, we used multilevel modelling (LeBreton & Senter, 2008). As a second step, the model was adjusted for the confounders (age, gender, education, and financial distress) (Model 2), with continuous variables centred. We did not control for mental health in our statistical models, as the multimorbidity variable already includes affective problems and chronic neurodegenerative diseases, such as Alzheimer’s disease. As the last step, we added our interest variable,

multimorbidity (Model 3). The deviance statistic is used to test if additional model predictors do improve the fit of the model. Odds ratios (OR), 95% confidence intervals (IC) and significance (where p-values of < 0.05 were considered statistically significant) are presented in the tables below. Statistical analyses were conducted using R software, version 4.0.2, and IBM SPSS Statistics 27.

RESULTS

In our sample, the mean age of the participants was 65.99 years (SD=9.43) and woman constituted 52.60 per cent of the sample. In addition, 31.30 per cent completed primary education or less, 40.61 per cent completed secondary education and 28.09 per cent completed post-secondary education. Moreover, across the study sample, 30.69 per cent of the respondents reported being financially distressed. Altogether 40.97 per cent of the participants reported the presence of multimorbidity, and 71.82 per cent of the respondents indicated having high precautionary behaviours.

Characteristics of the low and high precautionary behaviours groups are displayed in table 1. Among the respondents who reported multimorbidity, 72.2% engaged in high precautionary behaviours. Without controlling for confounders, all the variables listed in Table 1, except age, differed statistically in the two groups of individuals with high and low precautionary behaviours, although with no significant effect size.

Table 1 Characteristics of low and high precautionary behaviours groups


	low precautionary behaviours	high precautionary behaviours			Cohen's <i>d</i>	
	(N = 12930)	(N = 28602)	<i>t</i> / χ^2	p-value	/ phi	(CI 95%)
Age, mean(SD)	66.47 (9.75)	65.89 (9.29)	0.947	0.344	0.01	-0.011 - 0.031
Gender						
Male (%)	53.75	44.90	388.219	<0.001	0.10	0.087 - 0.106
Female (%)	46.26	55.10				
Education						
Primary or less (%)	24.9	33.83	151.751	<0.001	0.06	0.051 - 0.070
Secondary (%)	46.16	38.41				
Post-secondary (%)	28.94	27.76				
Financial distress						
No (%)	74.56	32.73	280.767	<0.001	0.08	0.072 - 0.092
Yes (%)	25.44	58.59				
Multimorbidity						
No (%)	60.14	58.59	116.211	<0.001	0.05	0.043 - 0.062
Yes (%)	39.86	41.41				

Source: Preliminary SHARE wave 8, release 0. Conclusions are preliminary. Weighted data, N=41534

Notes: t/χ^2 (t-test and chi-squared test), CI (confidence intervals). Tests for effect size: Cohen's *d*: small effect (≥ 0.20); medium effect (≥ 0.50); large effect (≥ 0.80); Phi: small effect (≥ 0.10); medium effect (≥ 0.30); large effect (≥ 0.50). Significant associations ($p < 0.05$) are in bold. The sample was limited to individuals aged 50+ who have left home since the outbreak of COVID-19.

Figure 1 shows the prevalence of high precautionary behaviours by country. Overall, the highest prevalence of precautionary behaviours was reported in Spain, Italy, Portugal, and Luxembourg (90.02; 89.49; 86.21; 85.17 per cent, respectively), while the lowest was reported in Sweden, Denmark, and Finland (32.59; 36.36; 43.64 per cent, respectively).

Fig. 1 Prevalence of high precautionary behaviours by country



Source: Preliminary SHARE, wave 8, release 0. Conclusions are preliminary.

Notes: *SE* Sweden; *DK* Denmark; *FI* Finland; *BG* Bulgaria; *LV* Latvia; *EE* Estonia; *SV* Slovakia; *CH* Switzerland; *DE* Germany; *IL* Israel; *CZ* Czech Republic; *ML* Malta; *HR* Croatia; *BE* Belgium; *CY* Cyprus; *FR* France; *HU* Hungary; *GR* Greece; *PL* Poland; *SI* Slovenia; *LT* Lithuania; *RO* Romania; *LU* Luxembourg; *PT* Portugal; *IT* Italia; *ES* Spain. Brackets denote 95% confidence intervals. The sample was limited to individuals aged 50+ who have left home since the outbreak of COVID-19.

The results of the multilevel logistic regression for precautionary behaviours are shown in Table 2. When Model 2 was compared with the null model, the deviance lowered, which means that, by adding the control variables, the model increased its quality (44221.4, p-value<0.001). When multimorbidity was included (Model 3), decreases in deviance were also observed (42717.2, p-value<0.001).

The final model (Model 3) showed that individuals with multimorbidity presented 25 per cent higher chances of having high precautionary behaviours compared with respondents without multimorbidity (OR = 1.25, 95 % CI [1.20;1.32]).

Women had 63 per cent more chances to engage in high precautionary behaviours compared to men (OR = 1.63, 95 % CI [1.55; 1.70]). Respondents with secondary and post-secondary education had more chances of adopting high precautionary behaviours, compared to respondents with primary education (OR= 1.08, 95 % CI [1.01; 1.15]; OR= 1.24, 95 % CI [1.16; 1.32], respectively). This is also the case for respondents who reported financial distress (OR = 1.11, 95 % CI [1.11; 1.25]) compared to their counterparts.

Table 2 Multilevel logistic regressions for precautionary behaviours

	Model 1	Model 2	Model 3
	OR (CI 95%)	OR (CI 95%)	OR (CI 95%)
Fixed parts			
(Intercept)	2.41 (1.80-3.21) ***	1.61 (1.21-2.15) **	1.45 (1.08-1.94)*
Age (years)		1.03 (1.01-1.06) **	1.01 (0.98-1.03)
Female		1.63 (1.56-1.71) ***	1.63 (1.55-1.70) ***
Education			
Primary or less		ref.	ref.
Secondary		1.07 (1.01-1.14) *	1.08 (1.01-1.15) *
Post-secondary		1.21 (1.13-1.29) ***	1.24 (1.16-1.32) ***
Financial distress		1.20 (1.13-1.27) ***	1.11 (1.11-1.25) ***
Multimorbidity			1.25 (1.20-1.32) ***
Random parts			
ICCcountry	0.136	-	-
Deviance	47809.9	44221.4	42717.2
N countries	26	26	26

Source: Preliminary SHARE, wave 8, release 0. Conclusions are preliminary.

Notes: *Ref* reference group, *OR* odds ratio, *CI* confidence intervals, *ICC* Intra-class Correlation Coefficients. Significant associations: '***' <0,001; '**' <0,01; '*' <0,05. The sample was limited to individuals aged 50+ who have left home since the outbreak of COVID-19.

DISCUSSION

Our results showed that individuals with multimorbidity were more likely to engage in precautionary behaviours than their counterparts. This result is consistent with the latest report from the English Longitudinal Study of Ageing (ELSA) COVID-19 Substudy, where individuals with multimorbidity had more precautionary behaviours, more specifically being isolated and staying at home (Zaninotto et al., 2020). In another recent study by Laires et al (2020), it was found that older Portuguese individuals with and without underlying health conditions had more self-awareness for the severity of COVID-19 and adopted more precautionary behaviours. Our results might be explained by theories of risk perception. Previous research into people's behaviours during pandemics concluded that the adoption of protective behaviours is consistent with these theories that point that people who perceive themselves as being more vulnerable to a threat tend to protect themselves more and engage in more precautionary behaviours (Bish & Michie, 2010; Rogers & Prentice-Dunn, 1997; Strecher et al., 1997; Wise et al., 2020). Therefore, self-awareness and perception of risk might explain why people with multimorbidity engage in more protective behaviours. Nevertheless, our conclusions differ from studies developed in the USA and Canada (O'Connor et al., 2020; Ramage-Morin & Polsky, 2020; Wolf et al., 2020), which showed that individuals with underlying health conditions were not engaging in more precautionary behaviours than their counterparts. Therefore, this topic should be addressed in further research.

Concerning gender, our results are in line with previous studies from authors who found that women adopted more protective behaviours against COVID-19, compared to men (Cvetković et al., 2020; Kim &

Crimmins, 2020; Lep et al., 2020; Lüdecke & von dem Knesebeck, 2020). The existing literature on risk perception can explain these results, as it states that women and men differ in their perception of risk (Gustafson, 1998).

Following the theories of risk perception, even though it would be expected that older adults would engage in more precautionary behaviours (Barber & Kim, 2020; Laires et al., 2020), our results showed no age differences between individuals with high and low precautionary behaviours.

Regarding education, our results are in line with studies that indicate that having a higher educational level is correlated with more precautionary behaviours (Bish & Michie, 2010; Zickfeld et al., 2020). Another interesting result is that the respondents who reported being more financially distressed had more chances of adopting more precautionary behaviours than those who did not indicate that they were financially distressed. We hypothesise that for individuals who are more financially distressed, it might be more difficult to afford medical assistance and medication, and therefore they adopted more measures to avoid becoming infected. This is, nevertheless, a research question to be investigated in the future.

In our descriptive analysis by country (figure 1), we found patterns according to geographical regions. In the Northern countries, such as Sweden, Denmark and Finland, the prevalence of precautionary behaviours was low. On the other hand, the prevalence was higher in Southern European countries, such as Spain, Italy and Portugal. Our hypothesis for these results lies in the prevalence of COVID-19 infection and mortality in Spain and Italy during the first wave of COVID-19 pandemic. Since these countries had a worse experience of the pandemic, their residents and those in neighbouring countries, like Portugal, were more likely to engage in precautionary behaviours because they may have perceived themselves as being more likely to get infected with COVID-19. Furthermore, the multiple health guidelines and restrictive measures imposed by the government may have been more effective in giving individuals perception of the severity of the risk. In contrast, during the first wave of COVID-19 pandemic, Northern Europe was not as severely affected, and so the guidelines and restrictive measures imposed were not as stringent, which gave individuals a weaker sense of risk perception.

Since the beginning of the pandemic, COVID-19 has caused increased morbidity and mortality around the globe, putting individuals with multimorbidity at a higher risk of significant harm. This study shows that public health messaging and actions must continue to be disseminated among middle and older aged individuals with multimorbidity to guarantee that this population continues to take preventive actions against COVID-19.

Additionally, more awareness campaigns should be aimed at middle and older aged men and individuals with less education and people with less financial distress, since these groups adopt fewer precautionary behaviours. The same is true for countries whose population adopts fewer precautionary behaviours, in order to make individuals aware of the risk, protect themselves and others and help mitigate the impact of the virus, regardless of their risk category.

Finally, yet importantly, with COVID-19, fast and large-scale behavioural changes are urgent and make it crucial for policymakers to be aware of whether the recommended guidelines are being followed or if there is a necessity for more awareness campaigns targeting people at greater risk of developing a more severe case of the disease. Thus, the results of this study are critical for obtaining a clear understanding of people's adoption of protective behaviours during the pandemic. This is essential for communication strategies and for addressing the present and future health crisis.

STRENGTH AND LIMITATIONS

To our knowledge, this is the first study analysing the association between multimorbidity and engaging in precautionary behaviours in middle and older aged Europeans in a cross-national perspective. Our sample is representative of the European and Israeli population, which allows us to perform generalizations. However, the findings of this research need to be interpreted within a framework sensitive to the limitations of the study. Since this study uses a cross-sectional design, we cannot assume causality. Furthermore, considering that obesity is now recognized as one of the main risk factors for COVID-19 severity, another limitation is the fact that we had no possibility of including this health condition in our study.

CONCLUSION

This study has several potential implications for middle and older aged individuals with multimorbidity, indicating that public health messages and guidelines should continue and should be reinforced to target this group as they are at a higher risk of developing worse health outcomes due to COVID-19. Since more pandemics may arise in the future, it is important to identify and target specific vulnerable groups in the early stages of an outbreak

in order to help contain and mitigate the spread, and also avoid increased morbidity and mortality. Public health safety messages should continue to be disseminated among the population, particularly in some countries.

REFERENCES

- Barber, S. J., & Kim, H. (2020). COVID-19 Worries and Behavior Changes in Older and Younger Men and Women. *The Journals of Gerontology: Series B, XX(Xx)*, 1–7. <https://doi.org/10.1093/geronb/gbaa068>
- Bergmann, M., Scherpenzeel, A., & Börsch-Supan, A. (2019). *SHARE Wave 7 Methodology: Panel Innovations and Life Histories* (M. Bergmann, A. Scherpenzeel, & A. Börsch-Supan (eds.)). Munich Center for the Economics of Aging (MEA). http://www.share-project.org/fileadmin/pdf_questionnaire_wave_7/w7_7.4.3.001_paper_questionnaire_main.pdf
- Bish, A., & Michie, S. (2010). Demographic and attitudinal determinants of protective behaviours during a pandemic: A review. *British Journal of Health Psychology*, *15*(4), 797–824. <https://doi.org/10.1348/135910710X485826>
- Boddington, N. L., Charlett, A., Elgohari, S., Walker, J. L., McDonald, H. I., Byers, C., Coughlan, L., Vilaplana, T. G., Whillock, R., Sinnathamby, M., Panagiotopoulos, N., Letley, L., MacDonald, P., Vivancos, R., Edeghere, O., Shingleton, J., Bennett, E., Grint, D. J., Strongman, H., ... Bernal, J. L. (2020). COVID-19 in Great Britain: Epidemiological and clinical characteristics of the first few hundred (FF100) cases: A descriptive case series and case control analysis. In *medRxiv* (p. 2020.05.18.20086157). medRxiv. <https://doi.org/10.1101/2020.05.18.20086157>
- Bonem, E. M., Ellsworth, P. C., & Gonzalez, R. (2015). Age Differences in Risk: Perceptions, Intentions and Domains. *Journal of Behavioral Decision Making*, *28*(4), 317–330. <https://doi.org/10.1002/bdm.1848>
- Börsch-Supan, A. (2020a). *Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 7. Release version: 7.1.0. SHARE-ERIC. Data set*. <https://doi.org/10.6103/SHARE.w7.710>
- Börsch-Supan A. (2020b). *Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 8. Release version: 0. SHARE-ERIC. Preliminary data set*
- Börsch-Supan, Axel, Brandt, M., Hunkler, C., Kneip, T., Korbmacher, J., Malter, F., Schaan, B., Stuck, S., & Zuber, S. (2013). Data resource profile: The survey of health, ageing and retirement in europe (share). *International Journal of Epidemiology*, *42*(4), 992–1001. <https://doi.org/10.1093/ije/dyt088>
- Cvetković, V. M., Nikolić, N., Nenadić, U. R., Öcal, A., Noji, E. K., & Zečević, M. (2020). Preparedness and preventive behaviors for a pandemic disaster caused by COVID-19 in Serbia. *International Journal of Environmental Research and Public Health*, *17*(11), 1–23. <https://doi.org/10.3390/ijerph17114124>
- Docherty, A. B., Harrison, E. M., Green, C. A., Hardwick, H. E., Pius, R., Norman, L., Holden, K. A., Read, J. M., Dondelinger, F., Carson, G., Merson, L., Lee, J., Plotkin, D., Sigfrid, L., Halpin, S., Jackson, C., Gamble, C., Horby, P. W., Nguyen-Van-Tam, J. S., ... Semple, M. G. (2020). Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: Prospective observational cohort study. *The BMJ*, *369*. <https://doi.org/10.1136/bmj.m1985>
- Ejaz, H., Alsrhani, A., Zafar, A., Javed, H., Junaid, K., Abdalla, A. E., Abosalif, K. O. A., Ahmed, Z., & Younas, S. (2020). COVID-19 and comorbidities: Deleterious impact on infected patients. *Journal of Infection and Public Health*, 1–7. <https://doi.org/10.1016/j.jiph.2020.07.014>
- Froes, M., Martins, B., & Neves, B. (2020). *C Omparison of M Ultimorbidity in Covid-19 Infected*.

- Glanz, K., & Bishop, D. B. (2010). The Role of Behavioral Science Theory in Development and Implementation of Public Health Interventions. *Annual Review of Public Health*, 31(1), 399–418. <https://doi.org/10.1146/annurev.publhealth.012809.103604>
- Guan, W. J., Liang, W. H., Zhao, Y., Liang, H. R., Chen, Z. S., Li, Y. M., Liu, X. Q., Chen, R. C., Tang, C. L., Wang, T., Ou, C. Q., Li, L., Chen, P. Y., Sang, L., Wang, W., Li, J. F., Li, C. C., Ou, L. M., Cheng, B., ... He, J. X. (2020). Comorbidity and its impact on 1,590 patients with Covid-19 in China: A nationwide analysis. *European Respiratory Journal*, 55(5). <https://doi.org/10.1183/13993003.00547-2020>
- Gustafson, P. E. (1998). Gender differences in risk perception: Theoretical and methodological perspectives. In *Risk Analysis* (Vol. 18, Issue 6, pp. 805–811). Plenum Publ Corp. <https://doi.org/10.1023/B:RIAN.0000005926.03250.c0>
- Iaccarino, G., Grassi, G., Borghi, C., Ferri, C., Salvetti, M., & Volpe Massimo, M. (2020). Age and multimorbidity predict death among COVID-19 Patients: Results of the SARS-RAS study of the Italian society of hypertension. *Hypertension*, 76(2), 1–7. <https://doi.org/10.1161/HYPERTENSIONAHA.120.15324>
- Jay, J., Bor, J., Nsoesie, E. O., Lipson, S. K., Jones, D. K., Galea, S., & Raifman, J. (2020). Neighbourhood income and physical distancing during the COVID-19 pandemic in the United States. *Nature Human Behaviour*. <https://doi.org/10.1038/s41562-020-00998-2>
- Jose, R., Narendran, M., Bindu, A., Beevi, N., L, M., & Benny, P. V. (2021). Public perception and preparedness for the pandemic COVID 19: A Health Belief Model approach. *Clinical Epidemiology and Global Health*, 9(June 2020), 41–46. <https://doi.org/10.1016/j.cegh.2020.06.009>
- Kim, J. K., & Crimmins, E. M. (2020). How does age affect personal and social reactions to COVID-19: Results from the national Understanding America Study. *PloS One*, 15(11), e0241950. <https://doi.org/10.1371/journal.pone.0241950>
- Laires, P. A., Dias, S., Gama, A., Moniz, A. M., Pedro, A. R., Soares, P., Aguiar, P., & Nunes, C. (2020). The Association of Chronic Diseases with COVID-19 Outcomes and its Role on Risk Perception: Nationwide COVID-19 Database & Online Community-Based Survey (Preprint). *JMIR Public Health and Surveillance*, 7, 1–12.
- LeBreton, J. M., & Senter, J. L. (2008). Answers to 20 Questions About Interrater Reliability and Interrater Agreement. *Organizational Research Methods*, 11(4), 815–852. <https://doi.org/10.1177/1094428106296642>
- Lep, Ž., Babnik, K., & Hacin Beyazoglu, K. (2020). Emotional Responses and Self-Protective Behavior Within Days of the COVID-19 Outbreak: The Promoting Role of Information Credibility. *Frontiers in Psychology*, 11, 1846. <https://doi.org/10.3389/fpsyg.2020.01846>
- Lüdecke, D., & von dem Knesebeck, O. (2020). Protective Behavior in Course of the COVID-19 Outbreak—Survey Results From Germany. *Frontiers in Public Health*, 8, 572561. <https://doi.org/10.3389/fpubh.2020.572561>
- Marengoni, A., Angleman, S., Melis, R., Mangialasche, F., Karp, A., Garmen, A., Meinow, B., & Fratiglioni, L. (2011). Aging with multimorbidity: A systematic review of the literature. *Ageing Research Reviews*, 10(4), 430–439. <https://doi.org/10.1016/j.arr.2011.03.003>
- O’Conor, R., Opsasnick, L., Benavente, J. Y., Russell, A. M., Wismer, G., Eifler, M., Marino, D., Curtis, L. M., Arvanitis, M., Lindquist, L., Persell, S. D., Bailey, S. C., & Wolf, M. S. (2020). Knowledge and Behaviors of Adults with Underlying Health Conditions During the Onset of the COVID-19 U.S. Outbreak: The Chicago COVID-19 Comorbidities Survey. *Journal of Community Health*, 45(6), 1149–1157. <https://doi.org/10.1007/s10900-020-00906-9>
- Ramage-Morin, P. L., & Polsky, J. Y. (2020). Health-related concerns and precautions during the COVID-19 pandemic: A comparison of Canadians with and without underlying health conditions. *Health Reports*, 31(5), 3–8. <https://doi.org/10.25318/82-003-x202000500001-eng>

- Rogers, R. W., & Prentice-Dunn, S. (1997). Protection motivation theory. In *Handbook of health behavior research 1: Personal and social determinants*. (pp. 113–132). Plenum Press.
- Sanyaolu, A., Okorie, C., Marinkovic, A., Patidar, R., Younis, K., Desai, P., Hosein, Z., Padda, I., Mangat, J., & Altaf, M. (2020). Comorbidity and its Impact on Patients with COVID-19. *SN Comprehensive Clinical Medicine*, 2(8), 1069–1076. <https://doi.org/10.1007/s42399-020-00363-4>
- Scherpenzeel, A., Axt, K., Bergmann, M., Douhou, S., Oepen, A., Sand, G., Schuller, K., Stuck, S., Wagner, M., & Börsch-Supan, A. (2020). Collecting survey data among the 50+ population during the COVID-19 pandemic: The Survey of Health, Ageing and Retirement in Europe (SHARE) SHARE-ERIC Central coordination Munich Center for the Economics of Aging (MEA), and Max Planck Institute for Social Law and Social Policy. *Survey Research Methods*, 14(2), 217–221. <https://doi.org/10.18148/srm/2020.v14i2.7738>
- Shankar, A., Saini, D., Roy, S., Jarrahi, A. M., Chakraborty, A., Bharati, S. J., & Taghizadeh-Hesary, F. (2020). Cancer care delivery challenges amidst coronavirus disease -19 (covid-19) outbreak: Specific precautions for cancer patients and cancer care providers to prevent spread. *Asian Pacific Journal of Cancer Prevention*, 21(3), 569–573. <https://doi.org/10.31557/APJCP.2020.21.3.569>
- Strecher, V. J., Champion, V. L., & Rosenstock, I. M. (1997). The health belief model and health behavior. In *Handbook of health behavior research 1: Personal and social determinants*. (pp. 71–91). Plenum Press.
- Tang, C. S. K., & Wong, C. Y. (2005). Psychosocial factors influencing the practice of preventive behaviors against the severe acute respiratory syndrome among older Chinese in Hong Kong. *Journal of Aging and Health*, 17(4), 490–506. <https://doi.org/10.1177/0898264305277966>
- Tang, & Wong. (2003). An Outbreak of the Severe Acute Respiratory Syndrome: Predictors of Health Behaviors and Effect of Community Prevention Measures in Hong Kong, China. *American Journal of Public Health*, 93(11), 1887–1889. <https://doi.org/10.2105/ajph.93.11.1887>
- Wise, T., Zbozinek, T. D., Michelini, G., Hagan, C. C., & Mobbs, D. (2020). Changes in risk perception and self-reported protective behaviour during the first week of the COVID-19 pandemic in the United States: COVID-19 risk perception and behavior. *Royal Society Open Science*, 7(9). <https://doi.org/10.1098/rsos.200742>
- Wolf, M. S., Serper, M., Opsasnick, L., O’Conor, R. M., Curtis, L., Benavente, J. Y., Wismer, G., Batio, S., Eifler, M., Zheng, P., Russell, A., Arvanitis, M., Ladner, D., Kwasny, M., Persell, S. D., Rowe, T., Linder, J. A., & Bailey, S. C. (2020). Awareness, Attitudes, and Actions Related to COVID-19 Among Adults With Chronic Conditions at the Onset of the U.S. Outbreak: A Cross-sectional Survey. *Annals of Internal Medicine*, 173(2), 100–109. <https://doi.org/10.7326/M20-1239>
- Zaninotto, P., Di Gessa, G., & Steel, N. (2020). The experience of older people with multimorbidity during the COVID-19 pandemic. *Elsa, September*.
- Zhang, J., Wang, X., Jia, X., Li, J., Hu, K., Chen, G., Wei, J., Gong, Z., Zhou, C., Yu, H., Yu, M., Lei, H., Cheng, F., Zhang, B., Xu, Y., Wang, G., & Dong, W. (2020). Risk factors for disease severity, unimprovement, and mortality in COVID-19 patients in Wuhan, China. *Clinical Microbiology and Infection*, 26(6), 767–772. <https://doi.org/10.1016/j.cmi.2020.04.012>
- Zickfeld, J. H., Schubert, T. W., Herting, A. K., Grahe, J., & Faasse, K. (2020). Correlates of Health-Protective Behavior During the Initial Days of the COVID-19 Outbreak in Norway. *Frontiers in Psychology*, 11(March), 1–19. <https://doi.org/10.3389/fpsyg.2020.564083>